

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

32-1312: IFN a 2a Recombinant Protein

Alternative Name: Leukocyte interferon, B cell interferon, Type I interferon, IFNA2, IFN-a 2a.

Description

Source: Escherichia Coli. Interferon Alpha Human 2a Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 165 amino acids and having a molecular mass of 19241 Dalton. The Interferon-a 2a gene was obtained from human leukocytes. The IFN-A 2a is purified by proprietary chromatographic techniques. IFN-alpha is produced by macrophages and has antiviral activities. Interferon stimulates the production of two enzymes: protein kinase and an oligoadenylate synthetase.

Product Info

Amount : 100 μg

Purification: Greater than 97.0% as determined by both:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: Lyophilized without additives.

Lyophilized Interferon alpha 2a although stable at room temperature for 3 weeks, should be

Storage condition: stored desiccated below -18°C. Upon reconstitution IFN-alpha 2a should be stored at 4°C

between 2-7 days and for future use below -18°C. For long term storage it is recommended to

add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids was determined and was found to be Cys-

Asp-Leu-Pro-Gln, conforming to the sequence of native human IFN-a.N-terminal methionine has

been completely removed enzymatically.

Application Note

It is recommended to reconstitute the lyophilized Interferon-alpha 2a in sterile $18M\tilde{A} \square \hat{A} \odot$ -cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. The specific activity as determined in a viral resistance assay using bovine kidney MDBK cells was found to be 270,000,000 IU/mg.

