

36-1849: Monoclonal Antibody to Myeloid-Associated Differentiation Marker (MYADM)(Clone : MYADM/971)

Clonality :	Monoclonal
Clone Name :	MYADM/971
Application :	FACS,IF,IHC
Reactivity :	Human
Gene :	MYADM
Gene ID :	91663
Uniprot ID :	Q96S97
Format :	Purified
Alternative Name :	MYADM,UNQ553/PRO1110
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant human MYADM protein

Description


Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This MAb is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Tonsil stained with MYDAM Monoclonal Antibody (MYADM/971).