

32-1360: mIL 1 alpha Recombinant Protein

Alternative Name : Hematopoietin-1,Lymphocyte-activating factor (LAF),Endogenous Pyrogen (EP),Leukocyte Endogenous Mediator (LEM),Mononuclear Cell Factor (MCF),IL-1 alpha,IL1,IL-1A,IL1F1.

Description

Source : Escherichia Coli. Interleukin-1A Mouse Recombinant produced in E.Coli is single, a non-glycosylated, Polypeptide chain containing 156 amino acids and having a molecular mass of 18 kDa. The IL-1A is purified by proprietary chromatographic techniques. Interleukin-1 alpha is a proinflammatory cytokine produced by a wide variety of cell types, including macrophages, osteoblasts, monocytes and hepatocytes. Circulating levels of are normally low and only rise after stimulation by agents such as those produced by inflammation, infection or microbial endotoxins. IL-1 alpha possesses a wide variety of biological activities and exerts its effects by binding to specific cell surface receptors.

Product Info

Amount : 10 µg
Purification : Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing 1X PBS, pH=7.4.
Storage condition : Lyophilized Interleukin-1a although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1A should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid : SAPYTYQSDL RYKLMKLVQR KFMNDSLNLQ TIYQDVVKHY LSTTWLNDLQ QEVKFDMYAY
SSGGDDSKYP VTLKISDSQL FVSAQGEDQP VLLKELPETP KLITGSETDL IFFWKSINSK NYFTSAAYPE
LFIATKEQSR VHLARGLPSP TDFQIS.

Application Note

It is recommended to reconstitute the lyophilized Interleukin 1a in sterile 18MΩ-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of murine D10S cells is < 0.002 ng/ml.

