

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1364: rmIL 1 alpha Recombinant Protein

Alternative Name Hematopoietin-1, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), IL-1 alpha, IL1, IL-1A, IL1F1.

Description

Source: Escherichia Coli. Recombinant IL 1 alpha Rhesus Macaque produced in E.coli cells is a non-glycosylated, homodimeric protein containing 159 amino acid chain and having a molecular mass of 18.1kDa. The IL 1 alpha is purified by proprietary chromatographic techniques. IL-1 alpha is produced by activated macrophages, stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1A proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Product Info

Amount:

Purification: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. Content: The IL 1 alpha was lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Lyophilized IL 1 alpha although stable at room temperature for 3 weeks, should be stored

desiccated below -18°C. Upon reconstitution IL 1 alpha should be stored at 4°C between 2-7 Storage condition:

days and for future use below -18°C.For long term storage it is recommended to add a carrier

protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: SAPFSFLSNM TYHFIRIIKH EFILNDTLNO TIIRANDOHL TAAAIHNLDE AVKFDMGAYT SSKDDTKVPV

ILRISKTQLY VSAQDEDQPV LLKEMPEINK TITGSETNFL FFWETHGTKN YFISVAHPNL FIATKHDNWV

CLAKGLPSIT DFOILENOA

Application Note

It is recommended to reconstitute the lyophilized IL 1 alpha Recombinant in sterile distilled H2O not less than 100Å∏µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by a cell proliferation assay using murine D10S cells is less than 10 pg/ml, corresponding to a specific activity of > 1.0x108IU/mg.

