

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1374: IL 1RA Recombinant Protein

Alternative Name: IRAP,IL1F3,IL1RA,IL-1ra3,ICIL-1RA,IL1RN,IL1 inhibitor,IL-1ra,MGC10430.

## **Description**

Source: Escherichia Coli. IL1ra Human Recombinant produced in E.Coli is a non-glycosylated, N-terminal methionyl form of the human naturally-occurring polypeptide chain containing 153 amino acids and having a molecular mass of 17000 Dalton. The IL1ra is purified by proprietary chromatographic techniques. Interleukin-1 ra is a member of the interleukin 1 cytokine family. This protein inhibits the activities of interleukin 1, alpha (IL1A) and interleukin 1, beta (IL1B), and modulates a variety of interleukin 1 related immune and inflammatory responses. This gene and five other closely related cytokine genes form a gene cluster spanning approximately 400 kb on chromosome 2. A polymorphism of this gene is reported to be associated with increased risk of osteoporotic fractures and gastric cancer. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.

## **Product Info**

Amount: 100 µg

**Purification:** Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: Lyophilized protein with no additives.

Lyophilized Interleukin 1ra although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-1ra should be stored at 4°C between 2-7 days Storage condition:

and for future use below -18°C.For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Amino Acid:** The sequence of the first five N-terminal amino acids was determined and was found to be Met-

Arg-Pro-Ser-Gly.

## **Application Note**

It is recommended to reconstitute the lyophilized Interleukin-1 ra in sterile 18M-cm H2O not less than 100̸Âμg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant inhibition of IL-1 stimulation of D10S cells was found to be 0.5 ng/ml corresponding to a Specific Activity of 2,000,000IU/mg.

