

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1384: IL 2 HEK Recombinant Protein(Discontinued)

Alternative Name: T-cell growth factor (TCGF), Aldesleukin, Lymphokine, IL-2.

Description

Source: HEK. Interleukin-2 Human Recombinant produced in HEK cells is a glycosylated monomer, having a total molecular weight of 15kDa. The IL2 is purified by proprietary chromatographic techniques. IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

Product Info

Amount: 10 µg

Purification: Greater than 95% as observeed by SDS-PAGE. **Content:** The IL2 was lyophilized from 1mg/ml in 1xPBS.

Lyophilized IL-2 although stable at room temperature for 3 weeks, should be stored desiccated

Storage condition : below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized IL-2 in sterile PBS containing 0.1% endotoxin-free recombinant HSA. The specific activity was determined by the dose-dependent stimulation of the proliferation of mouse CTLL-2 cells (mouse cytotoxic T cell line) and is typically 0.2-2ng/ml.

