

32-1389: qIL2 Recombinant Protein

Alternative Name : Interleukin-2,IL-2,T-cell growth factor,TCGF,IL2.

Description

Source : Escherichia Coli. Recombinant Equine Interleukin-2 produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 129 amino acids (with a substitution of S for C - at position 141 compared with the wild type IL2) and having a molecular mass of 14.9kDa.The IL-2 Equine is purified by proprietary chromatographic techniques. IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

Product Info

Amount :	20 µg
Purification :	Greater than 97.0% as determined by SDS-PAGE and HPLC analyses.
Content :	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 6.0.
Storage condition :	Lyophilized Interleukin-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	APTSSSKRET QQQLKQLQMD LKLLLEGVNN NKNPKLSKML TFKINMPKKA TELKHLQCLE EELKPLEEML KNFLSKDIKE LMSNINVTVL GLKGSETRFT CEYDDETGTI VEFLNKWITF SQSIFSTMT.

Application Note

It is recommended to reconstitute the lyophilized IL2 in sterile 18M-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by a cell proliferation assay using murine CTLL-2 cells is less than 1.0 Åµg/ml, corresponding to a specific activity of > 1000 IU/mg.

