

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1392: IL 3 Recombinant Protein

Alternative Name: MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

## **Description**

Source: Escherichia Coli. Interleukin-3 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 133 amino acids and having a molecular mass of 15000 Dalton. The IL-3 is purified by proprietary chromatographic techniques. IL3 is a potent growth promoting cytokine. This cytokine is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

## **Product Info**

Amount:  $10 \mu g$ 

**Purification:** Greater than 98.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE. **Content:** Lyophilized from a concentrated (1mg/ml) solution in water containing 0.3 mg/ml of NaHCO3.

Lyophilized Interleukin-3 although stable at room temperature for 3 weeks, should be stored

**Storage condition :** desiccated below -18°C. Upon reconstitution IL3 should be stored at 4°C between 2-7 days and

for future use below -18°C.Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids was determined and was found to be Ala-

Pro-Met-Thr-Gln.

## **Application Note**

It is recommended to reconstitute the lyophilized Interleukin-3 in sterile  $18M\tilde{A} \square \hat{A} \odot$ -cm H2O not less than  $100\tilde{A} \square \hat{A} \mu g/ml$ , which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of TF-1 cells is < 0.1 ng/ml, corresponding to a Specific Activity of 10,000,000 IU/mg.

