

32-1406: rmlL4 Recombinant Protein

Alternative Name : Interleukin-4,IL-4,B-cell stimulatory factor 1,BSF-1,Lymphocyte stimulatory factor 1,IL4.

Description

Source : Escherichia Coli. IL-4 Rhesus Macaque Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 129 amino acids and having a molecular mass of 14.9kDa.The IL4 Rhesus Macaque is purified by proprietary chromatographic techniques. IL4 is a pleiotropic cytokine produced by activated T cells. IL4 is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. IL4, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Product Info

Amount :	10 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.
Storage condition :	Lyophilized Interleukin-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL4 should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.
Amino Acid :	HNCHIALREI IETLSLSTEQ KTLCTKLIT DILAASKNTT EKETFCRAAT VLRQFYSHHE KDTRCLGATA QQFHRHKQLI RFLKRLDRNL WGLAGLNSCP VKEANQSTLE DFLERLKTIM REKYSKCSS.

Application Note

It is recommended to reconstitute the lyophilized Interleukin-4 in sterile 18MΩ-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependent stimulation of TF1 cells is less than 0.2ng/ml, corresponding to a specific activity of >5.0Å—106 IU/mg.

