

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1416: mIL 6 Recombinant Protein

Alternative Name : IFN-b2,B cell differentiation factor (BCDF),BSF-2,HPGF,HSF,MGI-2,IL-6,Interleukin HP-1,B-cell hybridoma growth factor.

Description

Source: Escherichia Coli. Interleukin-6 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 187 amino acids and having a molecular mass of 21709 Dalton. The IL-6 is purified by proprietary chromatographic techniques. Interleukin-6 is a potent pro-inflammatory cytokine primarily produced by activated T cells and an assortment of other cells including endothelial cells and macrophages. IL-6 affects B and T lymphocytes and has been shown to have a role in host defense, acute phase reactions, immune responses and hematopoiesis.

Product Info

Amount: $10 \mu g$

Purification: Greater than 96.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Content: Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Lyophilized Interleukin-6 although stable at room temperature for 3 weeks, should be stored

Storage condition : desiccated below -18°C. Upon reconstitution IL6 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: FPTSQVRRGD FTEDTTPNRP VYTTSQVGGL ITHVLWEIVE MRKELCNGNS DCMNNDDALA ENNLKLPEIQ

RNDGCYQTGY NQEICLLKIS SGLLEYHSYL EYMKNNLKDN KKDKARVLQR DTETLIHIFN QEVKDLHKIV

LPTPISNALL TDKLESQKEW LRTKTIQFIL KSLEEFLKVT LRSTRQT.

Application Note

It is recommended to reconstitute the lyophilized Mouse II-6 in sterile 18M-cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of the proliferation of IL-6-dependent murine 7TD1 cells is < 0.02 ng/ml, corresponding to a specific activity of > 50,000,000 units/mg.

