

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1427: IL 9 Recombinant Protein

Alternative Name: P40,HP40,T-cell growth factor p40,IL-9,P40 cytokine.

Description

Source: Escherichia Coli. Interleukin-9 Human Recombinant produced in E.Coli is a single, non-glycosylated single polypeptide chain containing 127 amino acids and having a molecular mass of 14,004 Dalton. The IL-9 is purified by proprietary chromatographic techniques. Factor that is thought to be a regulator of hematopoiesis. It has been shown to enhance the growth of human mast cells and megakaryoblastic leukemic cells as well as murine helper t-cell clones. IL-9 is a glycoprotein with a molecular weight of 32-39 that is derived from T-cells, and maps to human chromosome 5.

Product Info

Amount: 10 μg

Purification : Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. **Content :** Lyophilized from a concentrated (1mg/ml) solution containing 10mM Na2PO4, pH 7.5.

Lyophilized Interleukin-9 although stable at room temperature for 3 weeks, should be stored

desiccated below -18°C. Upon reconstitution IL9 should be stored at 4°C between 2-7 days and for

Storage condition:

desiccated below -18°C. Upon reconstitution it is should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1%)

HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: MQGCPTLAGI LDINFLINKM QEDPASKCHC SANVTSCLCL GIPSDNCTRP CFSERLSQMT

NTTMQTRYPL IFSRVKKSVE VLKNNKCPYF SCEQPCNQTT AGNALTFLKS LLEIFQKEKM

RGMRGKI.

Application Note

It is recommended to reconstitute the lyophilized Interleukin-9 in sterile 18MΩ-cm H2O not less than 100Âμg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependent stimulation of human MO7e cells is less than 0.2 ng/ml, corresponding to a Specific Activity of >5.0×106 IU/mg.

