

32-1435: rIL 10 Recombinant Protein

Alternative Name : B-TCGF,CSIF,TGIF,IL-10,IL10A,MGC126450,MGC126451,Cytokine synthesis inhibitory factor.

Description

Source : Escherichia Coli. IL-10 Recombinant Rat produced in E.coli is a single, glycosylated polypeptide chain containing 160 amino acids and having a molecular mass of 18.7 kDa. The Interleukin-10 Mouse is purified by proprietary chromatographic techniques. IL10 is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

Product Info

Amount :	10 µg
Purification :	Greater than 95.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein was lyophilized containing 25mM sodium phosphate buffer pH-7.25 and 100mM NaCl.
Storage condition :	Lyophilized IL-10 Rat although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Interleukin10 Rat recombinant should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	SKGHSIRGDN NCTHFPVSQT HMLRELRAAF SQVKTFQKK DQLDNILLTD SLLQDFKGYL GCQALSEMIK FYLVEVMPQA ENHGPEIKEH LNSLGEKLKT LWIQLRRCHR FLPCENKSKA VEQVKNDFNK LQDKGVYKAM NEFDIFINCI EAYVTLKMKN.

Application Note

It is recommended to reconstitute the lyophilized IL-10 Rat Recombinant in sterile 18MΩ·cm H₂O not less than 100 Åµg/ml, which can then be further diluted to other aqueous solutions. The ED₅₀ as determined by the dose dependent inhibition of MC/9 proliferation is less than 10ng/ml concentration corresponding to a Specific Activity of 100,000IU/mg.

