

## 32-1444: IL12 His Recombinant Protein

**Alternative Name** 23-219aa (IL12A) / 23-328aa (IL12B), Interleukin 12 (subunit beta/alpha), IL12 His, NKSF1, NKSF, CTL  
:  
maturation factor (TCMF), Cytotoxic lymphocyte maturation factor (CLMF), TSF, Edodekin-alpha, IL-12.

### Description

Source : Baculovirus. IL12 Human Recombinant produced in a baculovirus expression system is a glycosylated disulfide linked (through cysteines in bold) heterodimer comprised of IL12A (23-219aa, total of 203 aa, MW 23.3kDa) and IL12B (23-328aa, total of 306 aa, MW 34.6kDa), having a total predicted molecular mass of 57.9kDa (Molecular weight on SDS-PAGE will appear higher). IL12A is fused to a 6 amino acid His-Tag at C-terminus and purified by proprietary chromatographic techniques. IL-12 is a heterodimeric cytokine that stimulates the production of interferon gamma from T-cells and natural killer cells, and also induces differentiation of Th1 helper cells. IL-12 is an initiator of cell-mediated immunity.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 85.0% as determined by SDS-PAGE.
<b>Content :</b>	IL12 protein solution (0.2mg/ml) containing Phosphate buffered saline (pH7.4), 20% glycerol and 1mM EDTA.
<b>Storage condition :</b>	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
<b>Amino Acid :</b>	IL12A(p35):RNLVATPDP GMFPCLHHSQ NLLRAVS NML QKARQTLEFY PCTSEEIDHE DITKDKTSTV EACLPLELTK NESCLNSRET SFITNGSCLA SRKTSFMMAL CLSSYEDLK MYQVEFKTMN AKLLMDPKRQ IFLDQNMLAV IDELMQALNF NSETVPQKSS LEEPDFYKTK IKLCILLHAF RIRAVTIDRV MSYLNASHHH HHH. IL12B(p40):IWELKKDVYV VELD WYPDAP GEMVVLTCDT PEEDGITWTL DQSSEVLGSG KTLTIQVKEF GDAGQYTCHK GGEVLSHSL LLHKKEDGIW STDILKDQKE PKNKTF LRCE AKNYSGRFTC WWLTTISTDL TFSVKSSRGS SDPQGVTCGA ATLSAERV RG DNKEYEYSVE CQEDSACPAE EESLPIEVMV DAVHKLKYEN YTSSFFIRDI IKPDPPKNLQ LKPLKNSRQV EVSWEYPDTW STPHSYFSLT FCVQVQGKSK REKKDRVFTD KTSATVICRK NASISVRAQD RYYSSSWSEW ASVPCS.

### Application Note

The activity is determined by the IFN-g ELISA in a using NK-92 cell. The ED50 for this effect is less than or equal to 0.03 ng/ml.