

32-1463: mL 16 Recombinant Protein

Alternative Name : LCF,Lymphocyte Chemoattractant Factor,prIL-16,KIAA4048,mKIAA4048,IL16,IL-16,Interleukin-16.

Description

Source : Escherichia Coli. Interleukin-16 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 127 amino acids and having a molecular mass of 13.2 kDa. The Mouse IL-16 is purified by proprietary chromatographic techniques. IL-16 is a pleiotropic cytokine that functions as a chemoattractant, a modulator of T cell activation, and an inhibitor of HIV replication. The signaling process of IL-16 is mediated by CD4. The product of this gene undergoes proteolytic processing, which is found to yield two functional proteins. IL-16 functions exclusively attributed to the secreted C-terminal peptide, while the N-terminal product may play a role in cell cycle control. Caspase 3 is reported to be involved in the proteolytic processing of this protein. Two transcript variants encoding different isoforms have been found for this gene.IL-16 stimulates a migratory response in cd4+ lymphocytes, monocytes, and eosinophils. Also induces t-lymphocyte expression of interleukin 2 receptor. ligand for cd4.

Product Info

Amount :	10 µg
Purification :	Greater than 90.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis SDS-PAGE.
Content :	Murine IL-16 was lyophilized from 1mg/ml solution after extensive dialysis against 10mM sodium phosphate buffer, pH-7.5.
Storage condition :	Lyophilized Mouse IL-16 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution mouse IL16 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	MHDLNSSTDS AASASAASDI SVESKEATVC TVTLEKTSAG LGFSLEGGKG SLHGDKPLTI NRIFKGDRTG EMVQPGDEIL QLAGTAVQGL TRFEAWNVIK ALPDGPVTIV IRRSLQCKQ TTASADS.

Application Note

It is recommended to reconstitute the lyophilized Murine IL16 in sterile 18MΩ²Ω²-cm H2O not less than 100²Ω²µg/ml, which can then be further diluted to other aqueous solutions.