

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1475: mIL 17 Recombinant Protein

Alternative Name: CTLA-8,IL-17,IL-17A,Cytotoxic T-lymphocyte-associated antigen 8.

Description

Source: Escherichia Coli. Interleukin-17 Murine Recombinant produced in E.Coli is a homodimeric, non-glycosylated polypeptide chain containing a total of 268 (2x134 a.a.) amino acids and having a molecular mass of 30 kDa. The IL-17 is purified by proprietary chromatographic techniques. IL17 is a proinflammatory cytokine produced by activated T cells. IL-17 regulates the activities of NF-kappaB and mitogen-activated protein kinases. Interleukin-17 can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of IL-17 are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

Product Info

Amount: 25 µg

Purification: Greater than 98.0% as determined by:(a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Content: Lyophilized from a concentrated (1mg/ml) solution containing no additives.

Lyophilized Interleukin-17 although stable at room temperature for 3 weeks, should be stored

Storage condition: desiccated below -18°C. Upon reconstitution IL17 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1%)

HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: MAAIIPQSSA CPNTEAKDFL QNVKVNLKVF NSLGAKVSSR RPSDYLNRST SPWTLHRNED

PDRYPSVIWE AQCRHQRCVN AEGKLDHHMN SVLIQQEILV LKREPESCPF TFRVEKMLVG

VGCTCVASIV RQAA.

Application Note

It is recommended to reconstitute the lyophilized Interleukin 17 in sterile 18MΩ-cm H2O not less than 100Âμg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant induction of IL-6 production in cultured mouse NIH 3T3 fibroblasts was found to be 0.4 ng/ml, corresponding to a specific activity of 2,500,000 units/mg.

