

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1486: mIL 18 Recombinant Protein

Alternative Interferon-gamma-inducing factor,IGIF,IL-1g,IL-18,IL1F4,MGC12320,IFN-gamma-inducing

Name: factor,Interleukin-1 gamma,IL-1 gamma,Iboctadekin,Il18.

Description

Source: Escherichia Coli. Interleukin-18 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 178 amino acids (36-192 a.a.) and having a molecular mass of 20.4 kDa. The Mouse IL-18 is fused to a 20 amino acids His Tag at N-terminus and purified by proprietary chromatographic techniques. IL-18 is a proinflammatory cytokine. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

Product Info

Amount : 25 μg

Purification : Greater than 90.0% as determined by SDS-PAGE.

IL-18 is a proinflammatory cytokine. This cytokine can induce the IFN-gamma production of T

Content: cells. The combination of this cytokine and IL12 has been shown to inhibit IL4 dependent IgE

and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time.For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please avoid freeze thaw cycles.

Amino Acid: MGSSHHHHHH SSGLVPRGSH MNFGRLHCTT AVIRNINDQV LFVDKRQPVF EDMTDIDQSA SEPQTRLIIY

MYKDSEVRGL AVTLSVKDSK MSTLSCKNKI ISFEEMDPPE NIDDIQSDLI FFQKRVPGHN KMEFESSLYE

GHFLACQKED DAFKLILKKK DENGDKSVMF TLTNLHQS.

