

32-1582: LIF GST Recombinant Protein

Alternative Name : D factor, MLPLI, HILDA, Emfilermin, Leukemia Inhibitory factor, Differentiation-stimulating factor, Melanoma-derived LPL inhibitor.

Description

Source : Escherichia Coli. LIF produced in E.Coli is a single, non-glycosylated polypeptide chain containing 415 amino acids (23-202a.a.) and having a molecular mass of 47.2kDa. LIF is fused to a 236 amino acid His-GST tag at N-terminus & purified by proprietary chromatographic techniques. Leukemia inhibitory factor, is a pleiotropic cytokine which is expressed by numerous cells including activated T lymphocytes, monocytes, mast cells and neuronal cells. LIF takes part in the induction of hematopoietic differentiation in normal and myeloid leukemia cells, induction of neuronal cell differentiation, regulator of mesenchymal to epithelial conversion during kidney development, and is a key player in immune tolerance at the maternal-fetal interface.

Product Info

Amount : 10 µg
Purification : Greater than 90% as determined by SDS-PAGE.
Content : The LIF GST protein solution (1mg/1ml) is formulated in 20mM Tris-HCl buffer (pH 8.0) 1mM DTT, 50mM NaCl and 20% glycerol.
Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.
Amino Acid : MHHHHHHMSP ILGYWKIKGL VQPTRLLLEY LEEKYEEHLY ERDEGDKWRN KKFELGLEFP NLPYYIDGDV
 KLTQSMAIIR YIADKHNMLG GCPKERAIEIS MLEGAVLDIR YGVSRIAYSK DFETLKVDFL SKLPEMLKMF
 EDRLCHKTYL NGDHVTHPDF MLYDALDVVL YMDPMCLDAF PKLVCFFKRI EAIPQIDKYL KSSKYIAWPL
 QGWQATFGGG DHPPKSDLVP RGSMSPLPI TPVNATCAIR HPCHNNLMNQ IRSQLAQLNG SANALFILYY
 TAQGEPPNN LDKLCGPNTV DFPPFHANGT EKAKLVELYR IVVYLGTSLG NITRDQKILN PSALSLHSL
 NATADILRGL LSNVLCRLCS KYHVGHVDTV YGPDTSKDV FQKKKLGKQL LGKYKQIIAV LAQAF

