

## 32-1594: mM CSF Recombinant Protein

**Alternative Name :** CSF-1,Lanimostim,MCSF,M-CSF.

### Description

Source : Escherichia Coli. Macrophage Colony Stimulating Factor Mouse Recombinant produced in E.coli is a disulfide linked homodimer, non-glycosylated, polypeptide chain containing 2 x 156 amino acids and having a total molecular mass of 36.4 KD.MCSF is purified by proprietary chromatographic techniques. Granulocyte/Macrophage Colony-Stimulating Factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. CSF-1 induces cells of the monocyte/macrophage lineage. It plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
<b>Content :</b>	The protein was lyophilized with 0.5xPBS, pH 8.0.
<b>Storage condition :</b>	Lyophilized M-CSF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MCSF should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	MKEVSEHCSH MIGNGHLKVL QQLIDSQMET SCQIAFEFVD QEQLDDPVCY LKKAFFLVQD IIDETMRFKD NTPNANATER LQELSNLNS CFTKDYEEQN KACVRTFHET PLQLLEKIKN FFNETKNLLE KDOWNIFTKNC NNSFAKCSSR DVVTKP.

### Application Note

It is recommended to reconstitute the lyophilized M-CSF in sterile 18MÎ©-cm H2O not less than 100Âµg/ml, which can then be further diluted to other aqueous solutions. The ED50, as calculated by the dose-dependant stimulation of the proliferation of murine M-NFS-60 indicator cells is typically 1.1-1.6ng/ml corresponding to 909,090-625,000 U/mg.

