

32-1667: PEDF His Recombinant Protein

Alternative Name : Pigment epithelium-derived factor, PEDF, Serpin-F1, SerpinF1, EPC-1, EPC1, PIG35.

Description

Source : Escherichia Coli. PEDF Human Recombinant produced in E.Coli containing a natural variant M72T is a single, non-glycosylated, polypeptide chain containing 420 amino acids (20-418 a.a.) and having a total molecular mass of 46.7 kDa. PEDF is fused to a 20 amino acid His Tag at N-terminus and purified by proprietary chromatographic techniques. PEDF is a neurotrophic protein that induces extensive neuronal differentiation in retinoblastoma cells. SerpinF1 is a potent inhibitor of angiogenesis. EPC1 doesn't undergo the stressed to relaxed conformation transition characteristic as of the active serpins since it exhibits no serine protease inhibitory activity. Aqueous humour level of asymmetric dimethylarginine is correlated with PEDF in humans. ADMA and PEDF levels are increased in response to inflammation in uveitis. Lack of PEDF expression is a potent factor for the enhancement of tumor growth and angiogenesis in breast cancer. PEDF & VEGF genes contribute to the development of diabetic retinopathy. PEDF and VEGF structural changes in blood vessel wall play an important role in the pathophysiology of PD patients. PEDF-overexpressing tumors exhibited reduced intratumoral angiogenesis. SerpinF1 is a new promising approach for the treatment of osteosarcoma. Levels of the natural ocular anti-angiogenic factor SerturinF1 (PEDF) is associated with proliferative retinopathy. VEGF secreted by retinal pigment epithelial cells upregulates PEDF expression via VEGFR-1 in an autocrine manner. Serturin-F1 concentration in the aqueous humor of diabetic patients predicts who will develop progression of retinopathy. PEDF blocks angiogenic effects of leptin through its anti-oxidative properties.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µg |
| Purification : | Greater than 90% as determined by SDS-PAGE. |
| Content : | The PEDF solution contains 20mM Tris-HCl buffer(pH 8.0), 0.1M NaCl , and 20% glycerol. |
| Storage condition : | PEDF although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles. |
| Amino Acid : | MGSSHHHHHH SSGLVPRGSH MQNPASPPEE GSPDPDSTGA LVEEDPFFK VPVNKLAAAV SNFGYDLYRV RSSMSPTTNV LLSPLSVATA LSALSLGAEQ RTESIIHRAL YYDLISSPDI HGTYKELLD VTAPQKNLKS ASRIVFEKKL RIKSSFVAPL EKSYGTRPRV LTGNPRDLQ EINNWWQAQM KGKLARSTKE IPDEISILL GVAHFKGQWV TKFDSRKTSL EDFYLDEERT VRVPMMSDPK AVLRYGLDSD LSCKIAQLPL TGSMISIFFL PLKVTQNLTL IEESLTSEFI HDIDRELKTV QAVLTPKLK LSYEGEVTKS LQEMKLQSLF DSPDFSKITG KPIKLTQVEH RAGFEWNEDG AGTTPSPGLQ PAHLTFPLDY HLNQPFIFVL RDTDTGALLF IGKILDPRGP. |

