

## 32-1674: Pleiotrophin HEK Recombinant Protein(Discontinued)

**Alternative Name :**

PTN,Heparin Affin Regulatory Protein,HARP,Heparin-binding growth factor-8,HBGF-8,Osteoblast-Specific Factor-1,OSF-1,Heparin-binding growth-associated molecule,HB-GAM,HBNF-1 Heparin-binding brain mitogen,Heparin-binding neurite outgrowth-pro

### Description

Source : HEK. Pleiotrophin Human Recombinant produced in HEK cells is a non-glycosylated monomer, having a molecular weight of 18kDa.The Pleiotrophin is purified by proprietary chromatographic techniques. Pleiotrophin (Osteoblast-Specific Factor-1, OSF-1) contains 136 amino acid residues. The sequence is very rich in cationic amino acids (24% of the residues); lysine cluster sequences are found in the N-terminal and C-terminal ends of the structure.The OSF-1 gene was shown by Northern blotting analysis to be expressed in mouse calvarial osteoblast-enriched cells and in mouse brain tissues, but not in thymus, spleen, kidney, liver, lung, testis or heart. Pleiotrophin has the ability to promote adhesion, migration, expansion, and differentiation of human osteoprogenitor cells. In addition to certain types of cancer, the embryonic growth and differentiation factor pleiotrophin is found also in adults in inflammatory diseases. In osteoarthritis, pleiotrophin is especially expressed in early stages, and its concentrations in the synovial fluid could serve as a marker for the progress of the disease. Pleiotrophin might be involved in cartilage repair in osteoarthritis, in particular, in earlier stages.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 95% as observed by SDS-PAGE.
<b>Content :</b>	The Pleiotrophin was lyophilized from 1mg/ml in 1xPBS.
<b>Storage condition :</b>	Lyophilized Pleiotrophin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-7 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

### Application Note

It is recommended to reconstitute the lyophilized Pleiotrophin in sterile PBS containing 0.1% endotoxin-free recombinant HSA.