

32-1702: sCD40L Recombinant Protein

Alternative Name : CD40-L,Tumor necrosis factor ligand superfamily member 5,TNF-related activation protein,TRAP,T cell antigen Gp39,CD154 antigen,sCD40,IGM,IMD3,HIGM1,T-BAM,TNFSF5,hCD40L.

Description

Source : Escherichia Coli. sCD40 Human Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 149 amino acids and having a molecular mass of 16308 Dalton. The sCD40 is purified by proprietary chromatographic techniques. CD40L or CD154 is a membrane glycoprotein and differentiation antigen expressed on the surface of T-cells. The CD40 ligand stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines . CD40 ligand has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also costimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma , TNF-alpha , and IL2 .

Product Info

Amount : 50 µg
Purification : Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : Lyophilized from a 0.2µm filtered concentrated (1mg/ml) solution in PBS, pH 7.0.
Storage condition : Lyophilized sCD40 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CD154 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized sCD40 in sterile 18MΩ-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of IL-12 induction & IL-8 production by human PBMC was found to be 5-10 ng/ml.