

32-1721: sRANK Receptor Recombinant Protein

Alternative Name : Tumor necrosis factor receptor superfamily member 11A, Osteoclast differentiation factor receptor, ODFR, Receptor activator of NF- κ B, FEO, OFE, ODFR, OSTs, PDB2, RANK, CD265, OPTB7, TRANCER, LOH18CR1.

Description

Source : Escherichia Coli. sRANK Receptor Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 174 amino acids and having a molecular mass of 19.1kDa. The sRANK Receptor is purified by proprietary chromatographic techniques. sRANK Receptor is a part of the TNF superfamily of ligands and receptors which participates in the regulation of specific immunity and bone turnover. sRANK Receptor was originally acknowledged as a dendritic-cell-membrane protein, which by interacting with RANKL augments the capacity of dendritic cells to stimulate naive T cell proliferation and to endorse the survival of RANK and T cells. The full length human RANK cDNA encodes a type I transmembrane protein of 616 amino acids with a predicted 183 amino acid extracellular domain and a 383 amino acid cytoplasmic domain. sRANK Receptor is also expressed in a various tissues including skeletal muscle, thymus, liver, colon, small intestine and adrenal gland.

Product Info

Amount : 100 μ g
Purification : Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : Lyophilized from a 0.2 μ m filtered concentrated solution in 20mM Tris-HCl, pH 8.0 and 150mM NaCl.
Storage condition : Lyophilized sRANK Receptor although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution sRANK Receptor should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid : QIAPPCTSEK HYEHLGRCCN KCEPGKYMSS KCTTTSDSVC LPCGPDEYLD SWNEEDKCLL HKVCDTGKAL VAVVAGNSTT PRRCACTAGY HWSQDCECCR RNTECAPGLG AQHPLQLNKD TVCKPCLAGY FSDAFSSTDK CRPWTNCTFL GKRVEHHGTE KSDAVCSSSL PARK.

Application Note

It is recommended to reconstitute the lyophilized sRANK Receptor in sterile 18M-cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. The ED₅₀ as determined by its ability to inhibit sRANK Ligand induced nuclear factor kappa B (NF κ B) in RAW 264.7 cells is less than 50 ng/ml, corresponding to a specific activity of > 2.0 $\times 10^4$ IU/mg in the presence of 15 ng/ml of recombinant sRANK Ligand.

