

32-1723: mRELM a His Recombinant Protein

Alternative Name : Resistin-like alpha,RELMalpha,Cysteine-rich secreted protein FIZZ1,Parasite-induced macrophage novel gene 1 protein,Cysteine-rich secreted protein A12-gamma,RELM-a.

Description

Source : Escherichia Coli. RELM-alpha Mouse Recombinant is manufactured with a signal sequence of phage fd (20aa) and C-terminal fusion of flagTag (10aa). The RELM-alpha Flag-Tagged Fusion Protein is a 13.3 kDa protein containing 91 amino acid residues with 30 additional amino acid residues - signal sequence of phage fd, flagTag . Bronchoalveolar lavage fluid from mice with experimentally induced allergic pulmonary inflammation contains a novel 9.4 kDa cysteine-rich secreted protein, RELM-alpha (FIZZ1, found in inflammatory zone). RELM-alpha is a secreted protein that has a restricted tissue distribution with highest levels in adipose tissue stroma. Murine RELM-alpha (FIZZ1) is the founding member of a new gene family including two other murine genes expressed, respectively, in intestinal crypt epithelium (RELM-beta) and white adipose tissue (Resistin), and two related human genes.RELMalpha inhibits the differentiation of 3T3-L1 preadipocytes into adipocytes but has no effect on proliferation of 3T3-L1 preadipocytes. RELMalpha is able to form heterooligomers with resistin but not RELMbeta. Since RELMalpha is expressed by adipose tissue and it is a secreted factor, our findings suggest that RELMalpha may be involved in the control of the adipogenesis as well as in the process of muscle differentiation.In the lung, RELM-alpha is induced by hypoxia and was renamed as hypoxia-induced mitogenic factor (HIMF). HIMF strongly activated Akt phosphorylation. The phosphatidylinositol 3-kinase (PI3K) inhibitor LY294002 (10 micromol/L) inhibited HIMF-activated Akt phosphorylation. It also inhibited HIMFstimulated RPSM proliferation. Thus, the PI3K/Akt pathway, at least in part, mediates the proliferative effect of HIMF. Further studies showed that HIMF had angiogenic and vasoconstrictive properties. HIMF increased pulmonary arterial pressure and vascular resistance. Further studies suggest that HIMF regulates apoptosis and may participate in lung alveolarization and maturation.

Product Info

Amount :	25 µg
Purification :	Greater than 95% as determined by SDS-PAGE.
Content :	Filtered and lyophilized from 0.5 mg/ml in 5mM Tris pH 7.5, 25mM NaCl.
Storage condition :	Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid :	MKKLLFAIPL VVPFYSHSTM VNTDETIEII VENKVKELLA NPANYPSTVT TLSCTSVKT MNRWASCPAG MTATGCACGF ACGSWEIQSG DTCNCLCLLV DWTTARCCQL SLEDYKDDDD K.

Application Note

Add 0.2 ml of deionized H2O and let the lyophilized pellet dissolve completely.

