

## 32-1818: mTPO Recombinant Protein

**Alternative Name :** Megakaryocyte colony-stimulating factor, Myeloproliferative leukemia virus oncogene ligand, C-mpl ligand, ML, Megakaryocyte growth and development factor, MGDF, TPO, MKCSF, MPLLG, MGC163194, THPO.

### Description

Source : Escherichia Coli. Thrombopoietin Mouse Recombinant produced in E.Coli is a single, non-glycosylated soluble polypeptide chain containing 174 amino acids and having a molecular mass of 18704 Dalton. The TPO is purified by proprietary chromatographic techniques. Thrombopoietin is a glycoprotein hormone produced mainly by the liver and the kidney that regulates the production of platelets by the bone marrow. It stimulates the production and differentiation of megakaryocytes, the bone marrow cells that fragment into large numbers of platelets.

### Product Info

**Amount :** 10 µg  
**Purification :** Greater than 95.0% as determined by SDS-PAGE.  
**Content :** Lyophilized from a concentrated (1mg/ml) solution in water containing no additives.  
**Storage condition :** Lyophilized Thrombopoietin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TPO Mouse should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.  
**Amino Acid :** SPVAPACDPR LLNKLLRDSH LLHSRLSQCP DVDPLSIPVL LPAVDFSLGE WKTQTEQSKA QDILGAVSLL LEGVMAARGQ LEPSCLSSLL GQLSGQVRLL LGALQGLLGT QLPLQGRRTA HKDPNALFLS LQQLRGKVR FLLLVEGPTL CVRRTLPTTA VPSSTSQLLT LNKF.

### Application Note

It is recommended to reconstitute the lyophilized Thrombopoietin in sterile 18MΩ·cm H<sub>2</sub>O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions. ED50 range is less than 0.8ng/ml as determined by the dose dependent stimulation of Mo7e cells.

