

32-1891: GRO α , His Recombinant Protein

Alternative Name :

Growth-regulated protein alpha,CXCL1,Melanoma growth stimulatory activity,MGSA,Neutrophil-activating protein 3,NAP-3,GRO-alpha(1-73),chemokine (C-X-C motif) ligand 1,GRO1,GRO α ,SCYB1,MGSA-a,MGSA alpha.

Description

Source : Escherichia Coli. CXCL1 Human Recombinant produced in E.Coli is a single,non-glycosylated, polypeptide chain containing 94 amino acids (35-107) and having a molecular mass of 10.1 kDa. The CXCL1 is fused to a 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques. Chemokine (C-X-C motif) ligand 1 (CXCL1) is a small cytokine belonging to the CXC chemokine family that was previously called GRO1 oncogene, Neutrophil-activating protein 3 (NAP-3) and melanoma growth stimulating activity, alpha (MSGA-). It is secreted by human melanoma cells, has mitogenic properties and is implicated in melanoma pathogenesis. CXCL1 is expressed by macrophages, neutrophils and epithelial cells, and has neutrophil chemoattractant activity. CXCL1 plays a role in spinal cord development by inhibiting the migration of oligodendrocyte precursors and is involved in the processes of angiogenesis, inflammation, wound healing, and tumorigenesis. This chemokine elicits its effects by signaling through the chemokine receptor CXCR2. The gene for CXCL1 is located on human chromosome 4 amongst genes for other CXC chemokines.

Product Info

Amount :	50 μ g
Purification :	Greater than 90.0% as determined by SDS-PAGE.
Content :	CXCL1 in 20mM Tris-HCL pH-8 and 10% glycerol.
Storage condition :	Liquid CXCL1 although stable at 10°C for 1 week, should be stored below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	MGSSHHHHHH SSGLVPRGSH MASVATELRC QCLQTLQGIH PKNIQSVNVK SPGPHCAQTE VIATLKNRGR ACLNPAPIV KKIIEKMLNS DKS N.