w abeomics

32-1909: I 309 Recombinant Protein

Alternative Name : Small inducible cytokine A1,CCL1,T lymphocyte-secreted protein I-309,chemokine (C-C motif) ligand 1,P500,SISe,TCA3,I-309,SCYA1.

Description

Source : Escherichia Coli. I-309 Human Recombinant produced in E.Coli is a single,non-glycosylated, polypeptide chain containing 74 amino acids and having a molecular mass of 8504 Dalton. The I-309 is purified by proprietary chromatographic techniques. Chemokine (C-C motif) ligand 1 (CCL1) is a small glycoprotein secreted by activated T cells that belongs to a family inflammatory cytokines known as chemokines. CCL1 attracts monocytes, NK cells, and immature B cells and dendritic cells by interacting with a cell surface chemokine receptor called CCR8. This chemokine resides in a large cluster of CC chemokines on human chromosome 17.

Product Info

Amount : Purification :	10 μg Greater than 99.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The CCL1 protein was lyophilized with no additives.
Storage condition :	Lyophilized I-309 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	The sequence of the first five N-terminal amino acids was determined and was found to be Ser- Lys-Ser-Met-Gln.

Application Note

It is recommended to reconstitute the lyophilized I-309 in sterile $18M\tilde{A}$ \hat{a} cm H2O not less than $100\tilde{A}$ $\hat{a}\mu g/ml$, which can then be further diluted to other aqueous solutions. The Biological activity is calculated by its ability to chemoattract human T cells at 10-100ng/ml corresponding to a Specific Activity of 10,000-100,000IU/mg.

