

32-1922: rIP 10 Recombinant Protein

Alternative Name : C-X-C motif chemokine 10, 10 kDa interferon gamma-induced protein, Gamma-IP10, IP-10, Interferon-inducible protein 10, Protein Mob-1, Small-inducible cytokine B10, CXCL10, IIP10, Mob1, Scyb10.

Description

Source : Escherichia Coli. IP-10 Rat Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 77 amino acids and having a molecular mass of 8.6kDa. The IP-10 is purified by proprietary chromatographic techniques. Chemokine (C-X-C motif) ligand 10 (CXCL10) is a small cytokine belonging to the CXC chemokine family that is also known as 10 kDa interferon-gamma-induced protein (-IP10 or IP-10). CXCL10 is secreted by several cell types in response to IFN-. These cell types include monocytes, endothelial cells and fibroblasts. CXCL10 has been attributed to several roles, such as chemoattraction for monocytes and T cells, promotion of T cell adhesion to endothelial cells, antitumor activity, and inhibition of bone marrow colony formation and angiogenesis. The gene for CXCL10 is located on human chromosome 4 in a cluster among several other CXC chemokines. This chemokine elicits its effects by binding to the cell surface chemokine receptor CXCR3. The three-dimensional crystal structure of this chemokine has been determined under 3 different conditions to a resolution of up to 1.92Å.

Product Info

Amount : 25 µg
Purification : Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Content : Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.
Storage condition : Lyophilized IP-10 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL10 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid : IPLARTVRCT CIDFHEQPLR PRAIGKLEII PASLSCPHVE IIATMKKNNE KRCLNPESEA IKSLLKAVSQ RRSKRAP.

Application Note

It is recommended to reconstitute the lyophilized IP-10 in sterile 18MΩ·cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. Determined by its ability to chemoattract hCXCR3/HEK293 cells using a concentration range of 10.0-50.0 ng/ml.

