

32-1935: rMCP 1 Recombinant Protein

Alternative Name : Small inducible cytokine A2,CCL2,Monocyte chemotactic protein 1,MCP-1,Monocyte chemoattractant protein 1,Monocyte chemotactic and activating factor,MCAF,Monocyte secretory protein JE,HC11,chemokine (C-C motif) ligand 2,MCPI,SCYA2,GDCF-2,

Description

Source : Escherichia Coli. Monocyte Chemotactic Protein-1 Rat Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 125 amino acids and having a molecular mass of 14.1 kDa. The MCP-1 is purified by proprietary chromatographic techniques. Chemokine (C-C motif) ligand 2 (CCL2) is a small cytokine belonging to the CC chemokine family that is also known as monocyte chemotactic protein-1 (MCP-1). It is found at the site of tooth eruption and bone degradation. In the bone, CCL2 is expressed by mature osteoclasts and osteoblasts and is under the control of nuclear factor B (NFB). CCL2 recruits immune cells, such as monocytes, to sites of tissue injury and infection. This chemokine is produced as a protein precursor containing signal peptide of 23 amino acids and a mature peptide of 76 amino acids. It is a monomeric polypeptide, with a molecular weight of approximately 13kDa. As with many other CC chemokines, CCL2 is located on chromosome 17 in humans. The cell surface receptors that bind CCL2 are CCR2 and CCR5.

Product Info

Amount : 10 µg
Purification : Greater than 98.0% as determined by SDS-PAGE.
Content : The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives.
Storage condition : Lyophilized MCP-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid : QPDAVNAPLT CCYSFTGKMI PMSRLENYKR ITSSRCPEA VVFVTKLKRE ICADPNKEWV QKYIRKLDQN QVRSETTVFY KIASLRTSA PLNVNLTHKS EANASTLFST TTSSTSVEVT SMTEN.

Application Note

It is recommended to reconstitute the lyophilized Monocyte Chemotactic Protein-1 in sterile 18MΩ-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. ED₅₀ =1-10ng/ml corresponding to a Specific Activity of 100,000-1,000,000IU/mg. The biological activity was determined by measuring the dose dependent chemotaxis with human THP-1 cells. The optimal concentration should be determined for each specific application by an initial dose-response assay.