

## 32-1965: MIP 3b Recombinant Protein

### Alternative Name :

Small inducible cytokine A19,CCL19,Macrophage inflammatory protein 3 beta,MIP-3- beta,EBI1-ligand chemokine,ELC,Beta chemokine exodus-3,CK beta-11,chemokine (C-C motif) ligand 19,CKb11,MIP3B,MIP-3b,SCYA19,MGC34433.

### Description

Source : Escherichia Coli. Macrophage Inflammatory Protein-3 beta Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 77 amino acids and having a molecular mass of 8.8kDa. The MIP-3b is purified by proprietary chromatographic techniques. Chemokine (C-C motif) ligand 19 (CCL19) is a small cytokine belonging to the CC chemokine family that is also known as EBI1 ligand chemokine (ELC) and macrophage inflammatory protein-3-beta (MIP-3-beta). CCL19 is expressed abundantly in thymus and lymph nodes, with moderate levels in trachea and colon and low levels in stomach, small intestine, lung, kidney and spleen. The gene for CCL19 is located on human chromosome 9. This chemokine elicits its effects on its target cells by binding to the chemokine receptor chemokine receptor CCR7. It attracts certain cells of the immune system, including dendritic cells and antigen-engaged B cells.

### Product Info

**Amount :** 20 µg

**Purification :** Greater than 98.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

**Content :** Lyophilized without any additives.

**Storage condition :** Lyophilized MIP-3b although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL19 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

**Amino Acid :** GTNDAEDCCL SVTQKPIPGY IVRNHFYLLI KDGCRVPAVV FTTLRGRQLC APPDQPWVER IIQRLQRTSA KMKRRSS.

### Application Note

It is recommended to reconstitute the lyophilized Macrophage Inflammatory Protein-3b in sterile 18MΩ·cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The Activity of CCL19 is calculated by the ability to chemoattract Human T cells using a concentration of 7-40ng/ml corresponding to a Specific Activity of 25,000-143,000IU/mg.

