

32-1986: rmRantes Recombinant Protein

Alternative Name : Small inducible cytokine A5,CCL5,T-cell-specific RANTES protein,SIS-delta,T cell- specific protein P228,TCP228,chemokine (C-C motif) ligand 5,SISd,SCYA5,RANTES,D17S136E,MGC17164.

Description

Source : Escherichia Coli. Rantes Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 68 amino acids and having a molecular mass of 7.8kDa. The Mouse Rantes is purified by proprietary chromatographic techniques. Regulated upon Activation, Normal T-cell Expressed, and Secreted or RANTES is an 8 kDa protein classified as a chemotactic cytokine or chemokine. It has recently been renamed CCL5. RANTES is chemotactic for T cells, eosinophils and basophils and plays an active role in recruiting leukocytes into inflammatory sites. With the help of particular cytokines (i.e. IL-2 and IFN-) that are released by T cells, RANTES also induces the proliferation and activation of certain natural killer (NK) cells to form CHAK (CC-Chemokine-activated killer) cells. It is also a HIV-suppressive factor released from CD8+ T cells. This chemokine has been localized to chromosome 17 in humans.

Product Info

Amount : 20 µg
Purification : Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : Filtered (0.2µm) and lyophilized from a solution in 2xPBS, pH7.4.
Storage condition : Lyophilized Mouse Rantes although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Mouse CCL5 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid : SPYGSDDTTPC CFAYLSLALP RAHVKEYFYT SSKCSNLAVV FVTRNRQVC ANPEKKWVQE YINYLEMS.

Application Note

It is recommended to reconstitute the lyophilized Rantes in sterile 18MÎ©-cm H2O not less than 100Âµg/ml, which can then be further diluted to other aqueous solutions. The activity is determined by the ability to chemoattract total human lymphocytes population at a concentration between 1-10ng/ml.

