

## 32-1987: rRantes Recombinant Protein

**Alternative Name :** Small inducible cytokine A5,CCL5,T-cell-specific RANTES protein,SIS-delta,T cell- specific protein P228,TCP228,chemokine (C-C motif) ligand 5,SISd,SCYA5,RANTES,D17S136E,MGC17164.

### Description

Source : Escherichia Coli. Rantes Rat Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 68 amino acids and having a molecular mass of 7876 Dalton. The Rat Rantes is purified by proprietary chromatographic techniques. Regulated upon Activation, Normal T-cell Expressed, and Secreted or RANTES is an 8 kDa protein classified as a chemotactic cytokine or chemokine. It has recently been renamed CCL5. RANTES is chemotactic for T cells, eosinophils and basophils and plays an active role in recruiting leukocytes into inflammatory sites. With the help of particular cytokines (i.e. IL-2 and IFN-) that are released by T cells, RANTES also induces the proliferation and activation of certain natural killer (NK) cells to form CHAK (CC-Chemokine-activated killer) cells. It is also a HIV-suppressive factor released from CD8+ T cells. This chemokine has been localized to chromosome 17 in humans.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.  
**Content :** Lyophilized from a concentrated (1mg/ml) solution in water containing no additives.  
**Storage condition :** Lyophilized Rat Rantes although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Rat CCL5 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.  
**Amino Acid :** SPYGSDTTPC CFAYLSLALP RAHVKEYFYT SSKCSNLAVV FVTRRRNRQVC ANPEKKWVQE YINYLEMS.

### Application Note

It is recommended to reconstitute the lyophilized Rantes in sterile 18MΩ·cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The activity is determined by the ability to chemoattract total human lymphocytes and murine T-cells at a concentration between 1-10ng/ml corresponding to a Specific Activity of 100,000-1,000,000IU/mg.