

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-11108: Monoclonal Antibody to Cytokeratin, Basic (Type II or HMW) (Epithelial Marker)(Clone: SPM116)(Discontinued)

Clonality: Monoclonal SPM116

Application : FACS,IF,WB,IHC

Reactivity: Human, Monkey, Bovine, Dog, Rabbit, Mouse, Rat, Chicken

Gene: KRT76
Gene ID: 51350
Uniprot ID: Q01546
Format: Purified

Alternative Name : KRT76,KRT2B,KRT2P Isotype : Mouse IgG1 Kappa

Immunogen Information: Human epidermal keRatin

Description

This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. The acidic keratins have molecular weights (MW) of 56.5, 55, 51, 50, 50', 48, 46, 45, and 40kDa. This MAb recognizes the 65-67, 64, 59, 58, 56, and 52kDa keratins of basic subfamily. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. SPM115/SPM116 is a broad spectrum anti pan-keratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer.

Product Info

Amount : 100 μg

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly

toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.


Application Note

Flow Cytometry $(0.5-1\tilde{\mathbb{A}} \square \hat{\mathbb{A}} \mu g/million$ cells in 0.1ml); Immunofluorescence $(1-2\tilde{\mathbb{A}} \square \hat{\mathbb{A}} \mu g/ml)$; Western Blot $(0.5-1\tilde{\mathbb{A}} \square \hat{\mathbb{A}} \mu g/ml)$; Immunohistology (Formalin-fixed) $(0.25-0.5\tilde{\mathbb{A}} \square \hat{\mathbb{A}} \mu g/ml)$ for 30 min at RT); (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes); Optimal dilution for a specific application should be determined.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Cytokeratin, HMW Monoclonal Antibody (SPM116).