

14-126ACL: p53 LeeporTM Luciferase Reporter-HeLa Cell Line

Application : Functional Assay

Description

The p53 LeeporTM Luciferase Reporter cell line is a stably transfected HeLa cell line which expresses Renilla luciferase reporter gene under the transcriptional control of the p53 response element. p53 is a tumor suppressor that plays a crucial role in apoptosis and anticancer mechanisms. p53 reporter system is designed to monitor the p53-mediated signaling pathways. The p53 induction by doxorubicin is shown in Figure 1.

Product Info

Amount : 1 Vial
Content : Each vial contains $2 \sim 3 \times 10^6$ cells in 1 ml of 90% FBS + 10% DMSO.
Storage condition : Immediately upon receipt, store in liquid nitrogen.

Application Note

Application:

- Monitor the p53 induction activity.
- Screen for activators or inhibitors of the p53 signaling pathway.

Culture conditions:

Cells should be grown at 37°C with 5% CO₂ using DMEM medium (w/ L-Glutamine, 4.5g/L Glucose and Sodium Pyruvate) supplemented with 10% heat-inactivated FBS and 1% Pen/Strep, plus 3 µg/ml of Puromycin (Note: Puromycin can be omitted during the reporter cell assays).

It is recommended to quickly thaw the frozen cells upon receipt or from liquid nitrogen in a 37°C water-bath, transfer to a tube containing 10 ml of growth medium without Puromycin, spin down cells, resuspend cells in pre-warmed growth medium without Puromycin, transfer resuspended cells to T25 flask and culture in 37°C-CO₂ incubator.

Leave the T25 flask in the incubator for 1~3 days without disturbing or changing the medium until cells completely recover viability and become adherent. Once cells are over 90% adherent, remove growth medium and passage the cells through trypsinization and centrifugation. At first passage, switch to growth medium containing Puromycin. Cells should be split before they reach complete confluence.

To passage the cells, detach cells from culture vessel with Trypsin/EDTA, add complete growth medium and transfer to a tube, spin down cells, resuspend cells and seed appropriate aliquots of cells suspension into new culture vessels. Subcultivation ration = 1:10 to 1:20 weekly. To achieve satisfactory results, cells should not be passaged over 16 times.

Functional validation:

A. Response of p53 LeeporTM – HeLa cells to doxorubicin.

