

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-4171-F: Monoclonal antibody to CD68 (Clone: ABM53F5)-FITC Conjugated

Clonality: Monoclonal **Clone Name:** ABM53F5 Application: FACS Reactivity: Human Conjugate: FITC Gene: **CD68** Gene ID: 968 **Uniprot ID:** P34810 Format: **Purified**

Alternative Name : Macrosialin, Gp110 **Isotype :** mouse IgG2b, k

Immunogen Information: A partial length recombinant protein (a.a 80-280) of CD68 was used as the immunogen for

this antibody.

Description

CD68 is a transmembrane glycoprotein which acts as a commercialized marker for human monocytes and macrophages. Its expression is thought to be regulated by a macrophage-specific promoter. However, several immunohistochemical studies have indicated that CD68 antibodies also react with other haematopoietic and non-haematopoietic cell types. It also can stain myeloid cells, dendritic cells, fibroblasts, Langerhans cells and others. In human monocytes, which produce high levels of CD68 mRNA, the gene is characterized by intramolecular ligations between the promoter and the 3' intervening region. CD68 is also used as a pan-macrophage marker for TAM (Tumor-Associated Macrophages) which always involve in carcinogenesis.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 0.2 mg/ml in Tris buffer containing 0.05% Azide **Storage condition:** Store the antibody at 4°C; stable for 6 months.

Application Note

FACS: 0.5-1 μg/10⁶ cells



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig:1- Cell surface Flow analysis of hCD68 in human PBMC (Monocytes) using 0.5 μ g/ 10^6 cells. Green represents isotype control (ABEOMICS);red represents anti-hCD68 FITC conjugated antibody (10-4171F).