

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-3021-AC: Monoclonal Antibody to TLR4 (Clone: HTA125) APC Conjugated

Clonality: Monoclonal Clone Name: HTA125 Application: **FACS** Reactivity: Human Conjugate: APC Gene: TLR4 Gene ID: 7099 **Uniprot ID:** O00206 Format: Purified TLR4 **Alternative Name:** 

**Isotype:** Mouse IgG2a, kappa

Immunogen Information: This antibody was developed by immunizing mice with Ba/F3 cell line expressing human TLR4 cell

surface antigen.

## **Description**

TLR4 (Toll-like receptor-4) is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. It primarily mediates cellular signaling induced by Gram negative bacteria. TLR4 is highly specific for LPS and it is associated with CD14, a co-receptor for LPS. The downstream signaling pathway of TLR4 leads to the activation of NF-kappaB through myeloid differentiation protein (MyD88) and IL-1 receptor-associated kinase in various cell types. Keratinocytes express TLR4 at the mRNA and protein levels whereas TLR4 are also present in the normal human epidermis in vivo and their expression is regulated by microbial components. TLR4 is expressed on CD4+ T cells as well, the functional significance of which is unclear.

## **Product Info**

Amount: 25 Test / 100 Test

**Purification:** Protein G Chromatography

Content: 0.04 mg/ml in PBS with 0.05% Azide.

**Storage condition :** Store the antibody at 4°C, stable for 6 months.

## **Application Note**

FACS analysis: 0.5-1 µg/10^6



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-1:Intracellular flow analysis of hTLR4 APC conjugated antibody in hPBMC (Monocytes) using 0.5  $\mu$ g/10^6 cells of hTLR4 antibody (Clone: HTA125). Green represents isotype control; red represents anti-hTLR4 APC conjugated antibody.