

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

11-8018: Polyclonal Antibody to LST8/GBL

Clonality: Polyclonal

Application: WB
Reactivity: Human
Gene: MLST8
Gene ID: 64223
Uniprot ID: Q9BVC4
Format: Purified

Alternative Name: MLST8,GBL,LST8
Isotype: Rabbit IgG

Immunogen Information: A partial length recombinant LST8/GBL protein (amino acids 100-320) was used as the

immunogen for this antibody.

Description

MLST8 or GbL (G protein beta protein subunit-like) is a subunit of mTORC1 and mTORC2 which regulates cell growth and survival. Both mTORC1 and 2 are activated in response to growth factors or amino acids but mTORC2 is most likely insensitive towards nutrients. Within mTORC1, LST8 interacts directly with mTOR and helps to enhance its kinase activity. mTORC1 phosphorylates EIF4EBP1 and thereby releasing it from inhibiting the elongation initiation factor 4E (eiF4E). mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. LST8/GBL more or less ubiquitously expressed with highest levels in skeletal muscle, heart and kidney.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein A Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 1-3 ̸µg/ml.




Fig-1: Western blot analysis of LST8/GBL. Anti-LST8/GBL (11-8018) was used at 1 μ g/ml on Kidney lysate.