

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

11-7005: Polyclonal Antibody to DMAP1

Clonality: Polyclonal

Application: WB
Reactivity: Human
Gene: DMAP1
Gene ID: 55929
Uniprot ID: Q9NPF5
Format: Purified

Alternative Name: DMAP1,KIAA1425

Isotype: Rabbit IgG

Immunogen Information: A partial length recombinant DMAP1 protein (amino acids 140-462) was used as the immunogen

for this antibody.

Description

DNA methyltransferase 1-associated protein 1 is an enzyme that in humans is encoded by the DMAP1 gene and consisting of 433 aminoacids. It is involved in transcription repression and activation. DMAP1 interacts with HDAC2 which provides a mechanism for histone deacetylation in heterochromatin following replication of DNA at late dismissal origins. DMAP1 can also repress transcription independently of histone deacetylase activity. It participates in the nuclear localization of URI1 and increases its transcriptional co-repressor activity. It is expressed in adipose tissue, adrenal gland, amniotic fluid, cerebellar cortex, hippocampus, liver, etc.

Product Info

Amount : 25 μg / 100 μg

Purification: Protein A Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 µg/ml




Fig-1: Western blot analysis of DMAP1. Anti-DMAP1 antibody (11-7005) was used at 4 μ g/ml on 1) Hela, 2) HepG2 lysates.