

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 11-7007: Polyclonal Antibody to Histone H3.3

Clonality: Polyclonal
Application: IHC,WB
Reactivity: Mouse,Human

**Gene :** H3F3A **Gene ID :** 3020

Uniprot ID: P84243
Format: Purified

Alternative Name: H3F3A,H3.3A,H3F3,PP781

**Isotype:** Rabbit IgG

Immunogen Information: Full length recombinant Histone H3.3 protein was used as the immunogen for this antibody.

## **Description**

Histones are major chromatin proteins involved in nucleosome assembly. Their major function is to compact long DNA strands inside the nucleus and chromatin regulation. Histone H3.3 is a replication independent variant of Histone H3 which is generally expressed in quiescent or terminally differentiated cells. Modifications in Histone H3.3 are associated with transcriptional activation and acts as an epigenetic mark of transcriptionally active chromatin. Histone H3.3 is a 15 kDa protein which is expressed throughout the cell cycle independently of DNA synthesis in all eukaryotic genomes.

## **Product Info**

**Amount :**  $25 \mu g / 100 \mu g$ 

**Purification:** Protein A Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 1-2  $\mu$ g/ml, Immunohistochemical analysis: 5  $\mu$ g/ml




Figure-1: Western blot analysis of Histone H3.3. Anti-Histone H3.3 antibody (11-7007) was used at 2  $\mu$ g/ml on (1) Raji, 2) Jurkat lysates.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2: Immunohistochemical analysis of Histone H3.3. Anti-Histone H3.3 antibody (11-7007) in human Lung tissue at 5  $\mu$ g/ml.




Figure-3: Western blot analysis of Histone H3.3. Anti-Histone H3.3 antibody (11-7007) was used at 2  $\mu$ g/ml on (1) K562, 2) RAW, (3) EL-4 and (4) m Testis lysates.