

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-3003: Monoclonal Antibody to TLR9 (Clone: ABM1C51)

Clonality: Monoclonal Clone Name: ABM1C51 Application: IHC,FACS,WB Reactivity: Mouse, Human

Gene: TLR9 Gene ID: 54106 **Uniprot ID: Q9NR96** Format: Purified

**Alternative Name:** TLR9,UNQ5798/PRO19605

Isotype: Mouse IgG1 Kappa

A partial length recombinant TLR9 protein (amino acids 100-290) was used as the immunogen for **Immunogen Information:** 

the antibody.

## **Description**

TLR9, a member of toll-like receptor family are central to the innate immunity by identifying pathogen associated molecular patterns (PAMPs). TLR9 identify unmethylated CpG dinucleotides present in bacterial DNA leading to NF-kB activation.

## **Product Info**

Amount:  $25 \mu g / 100 \mu g$ 

Purification: Protein G Chromatography

25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium Content:

azide is highly toxic.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 2-4 µg/ml, FACS analysis: 0.5 µg/10^6 cells, Immunohistochemical analysis: 5 µg/ml




Fig-1: Immunohistochemical analysis of TLR9 in human stomach tissue using TLR9 antibody (Clone: ABM1C51) at 5 µg/ml.



Email: info@abeomics.com






Fig-2: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 2  $\mu$ g/ml on (1) Raji and (2) EL4 lysates.




Fig-3: Intracellular flow analysis of TLR9 in human PBMC (Lymphocytes) using 0.5 μg/10^6 cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary.




Fig-4: Intracellular flow analysis of TLR9 in Raji cells using  $0.5 \mu g/10^6$  cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Fig-5 : Immunohistochemical analysis of TLR9 in human Liver tissue using TLR9 antibody (Clone: ABM1C51) at 20  $\mu$ g/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-6: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 4  $\mu$ g/ml on Daudi lysate.