

## 10-1052: Monoclonal Antibody to Bcl-2 (Clone: BC1)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	BC1
<b>Application :</b>	IHC,FACS,WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	BCL2
<b>Gene ID :</b>	596
<b>Uniprot ID :</b>	P10415
<b>Format :</b>	Purified
<b>Alternative Name :</b>	BCL2
<b>Isotype :</b>	Mouse IgG1 Kappa
<b>Immunogen Information :</b>	A synthetic peptide corresponding to amino acids 41-54 (GAAPAPGIFSSQPG-Cys), of human Bcl-2 was used as immunogen.

### Description

Bcl-2 (B-cell lymphoma 2), is a member of Bcl-2 family of regulator proteins. These proteins contain a hydrophobic cleft that binds to BH3-only proteins and to the pro-apoptotic Bcl-2 family members Bad, Bak, and Bax to inhibit apoptosis. In the absence of this binding, the proapoptotic Bcl-2 members are recruited to the OMM (Outer Mitochondrial Membrane) at which they oligomerize and cause OMM permeabilization, releasing proapoptotic effectors such as SMAC or cytochrome-c. Bcl-2 also neutralize a group of 'sensor' proteins (such as BIM), which are triggered by cytotoxic stimuli such as chemotherapy. BCL-2 proteins therefore have a central role as guardians against apoptosis, helping cancer cells to evade cell death.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Western blot analysis: 2-4 µg/ml, Immunohistochemical analysis: 5 µg/ml, Flowcytometric analysis- 2-4 µg/10<sup>6</sup> Cells




Figure-1: Western blot analysis of Bcl2. Anti- Bcl2(Clone: BC1) was used at 2 µg/ml on 293, MCF7 and Jurkat lysates.




Figure-2 : Immunohistochemical analysis of Bcl-2 in human kidney tissue using Bcl-2 antibody (Clone: BC1) at 5 µg/ml.




Figure-3 : Immunohistochemical analysis of Bcl-2 in human Prostate tissue using Bcl-2 antibody (Clone: BC1) at 5 µg/ml.




Figure-4 : Immunohistochemical analysis of Bcl-2 in human Spleen tissue using Bcl-2 antibody (Clone: BC1) at 5 µg/ml.




Figure-5 : Immunohistochemical analysis of Bcl-2 in human Stomach tissue using Bcl-2 antibody (Clone: BC1) at 5 µg/ml.




Figure-6: Intracellular flow cytometric analysis of Bcl-2 in Jurkat cell lines using 2 µg/10<sup>6</sup> cells of Anti-Bcl-2 antibody (10-1052 Abeomics) . Green represent isotype control and red represent Anti-Bcl-2 antibody (Clone: BC1). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.