

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

20-1051: Polyclonal antibody to Caspase-10/FLICE2

 Clonality :
 Polyclonal

 Application :
 WB,IHC,IP

 Reactivity :
 Human

 Gene :
 CASP10

 Gene ID :
 843

 Uniprot ID :
 Q92851

 Format :
 Sera

Alternative Name : CASP10,MCH4
Isotype : Rabbit IgG

Immunogen Information: A recombinant full-length human Caspase-10/FLICE2 protein was used as the immunogen for this

antibody

Description

The current dogma of apoptosis suggests that the components of the core cell-death machinery are integral to cells and widely conserved across species. Caspases are typically divided into 3 major groups, depending on the structure of their prodomain and their function. Group 1: inflammatory caspases (caspases 1, 4, 5, 11, 12, 14). Group II: initiator of apoptosis caspases (caspases 2, 8, 9). Group III: effector caspases (caspases 3, 6, 7). Caspases are constitutively expressed in almost all cell types as inactive proenzymes (zymogens: enzyme precursors which require a biochemical change to become active enzymes) that are processed and activated in response to a variety of pro-apoptotic or inflammatory stimuli. The procaspases (32-56 kDa) contain four domains: an N-terminal prodomain (2-25 kDa), a large subunit (p20: 17-21 kDa), a small subunit (p10: 10-13 kDa) and a short linker region between the large and small subunits. Caspase activation involves proteolytic processing of the proenzyme at specific aspartate residues between the domains. This results in removal of the prodomain as well as the linker region and formation of a heterodimer containing one large and one small subunit (p20-p10). However, caspase-10 produces, by alternative splicing, at least 12 different transcripts, all with introns, putatively encoding at least 11 different protein isoforms.

Product Info

Amount : $50 \mu l$ Content : $50 \mu l$ sera

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200