

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 20-1066: Polyclonal antibody to CARD12 (IPAF/CIAN)

Clonality: Polyclonal
Application: WB,IHC,IP
Reactivity: Human,Mouse

Gene: NLRC4
Gene ID: 58484
Uniprot ID: Q9NPP4
Format: Sera

Alternative Name: NLRC4,CARD12,CLAN,CLAN1,IPAF,UNQ6189/PRO20215

**Isotype:** Rabbit IgG

Immunogen Information: A synthetic peptide of IPAF/CIAN/CARD12 (amino acis 637-654 EEAPETYIPSRAVSLFFN) was

used as immunogen for this antibody

## **Description**

Ipaf (also known as Clan/CARD12) is a CARD domain containing protein. In general CARD proteins are implicated in host defense against infection, environmental stress or cellular damage. CARD domains are found in the N-terminal pro-domains of certain caspases, a family of apoptotic and pro-inflammatory proteases, as well as in a diversity of other proteins including Ipaf/Clan/CARD12. There are at least three major signaling pathways in which CARD proteins act: (1) Regulation of caspase activation in the context of apoptosis (2) Regulation of caspase activation in the context of inflammation (3) Regaultion of NF-kB activation in the context of innate or adaptive immune responses. As there is significant crosstalk between pathways that lead to caspase-mediated apoptosis or inflammation and pathways that result in NF-kB activation, it is logical that similar protein modules such as CARD domains are found repeatedly in proteins from all three pathways. Ipaf plays a role in regulating caspase-1 activity, which in turn mediates the maturation of inflammatory cytokines IL-1b and IL-18. Ipaf also interact with the pro-apoptotic adaptor protein ASC and co-expression of Ipaf with ASC has been shown to induce NF-kB activation and apoptosis.

## **Product Info**

Amount :  $50 \mu l$  Content :  $50 \mu l$  sera

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200