

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

20-1095: Polyclonal antibody to TRAF-2

Clonality: Polyclonal Application: WB,IHC,IP

Reactivity: Human, Mouse, Rat

 Gene :
 TRAF2

 Gene ID :
 7186

 Uniprot ID :
 Q12933

 Format :
 Sera

Alternative Name: TRAF2,TRAP3 Isotype: Rabbit IgG

Immunogen Information: A synthetic peptide of human TRAF-2 (amino acids 87-106 SSSAFPDNAARREVESLPAV) was

used as immunogen for this antibody

Description

This antibody recognizes TRAF2, which is a member of TRAF (TNF receptor-associated factor) adapter proteins composed 501 amino acids. It Links TNFRs (TNF Receptors) to the SAPKs (Stress-Activated Protein Kinases) and p38s. TRAF2 can activate ASK1 (Apoptosis Signal-Regulating Kinase-1) in vivo and can interact in vivo with the amino- and carboxyl-terminal noncatalytic domains of the ASK1 polypeptide. Expression of the amino-terminal noncatalytic domain of ASK1 can inhibit TNF and TRAF2 activation of SAPK. TRAF2 is a potential mediator of CD40 signaling. In vitro, TRAF2 and TRAF3 bind to the CD40 Cytoplasmic Tail (CD40CT) with much higher affinity than TRAF5 and TRAF6 (TNF Receptor Associated Factor-6) and that TRAF2 and TRAF3 bind to different residues of the CD40CT. The TRAF2-binding site of the CD40CT is critical for NF-KappaB and SAPK activation, as well as the UP-regulation of the ICAM1 (Intercellular Adhesion Molecule-1).

Product Info

Amount : $50 \mu l$ Content : $50 \mu l$ sera

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200




Fig:1 Immunhistochemical analysis of TRAF2 in formalin-fixed, paraffin-embedded normal human thymus sections using 20-1095 at 1:2000. A-C, successively higher magnifications. Hematoxylin-eosin counterstain.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig:2 Immunhistochemical analysis of TRAF2 in formalin-fixed, paraffin-embedded normal human bone marrow section using 20-1095 at 1:2000. Hematoxylin-eosin counterstain.