

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-7564: Monoclonal Antibody to FLT1 (Clone: ABM4A56)

Clonality: Monoclonal ABM4A56 Clone Name: Application: IHC Reactivity: Human Gene: FLT1 Gene ID: 2321 **Uniprot ID:** P17948 Format:

**Alternative Name:** FLT1,FLT,FRT,VEGFR1 Isotype: Mouse IgG2b Kappa

A partial length recombinant FLT1 protein (amino acids 102-321) was used as the immunogen for **Immunogen Information:** 

this antibody.

Purified

## **Description**

FLT-1 (fms-related tyrosine kinase 1) is an endogenous inhibitor of vascular endothelial growth factor, which is involved in cardiovascular remodeling and atherosclerosis development. It potently inhibits angiogenesis by binding extracellularly to VEGF (vascular endothelial growth factor). FLT-1 is expressed on tumor cells and is implicated in tumor growth and progression. It is overexpressed in primary tumors and nodal metastasis. FLT-1 regulates monocyte migration, recruits endothelial cell progenitors, increases the adhesive properties of natural killer cells and induces growth factors. It is observed in many diseases with endothelial dysfunction, including pre-eclampsia and diabetic retinopathy, but whether it contributes to endothelial injury in IgAN requires further exploration.

## **Product Info**

Amount:  $25 \mu g / 100 \mu g$ 

Purification: Protein G Chromatography

25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium Content:

azide is highly toxic.

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

Immunohistochemical analysis: 5-10 µg/ml




Fig-1: Immunohistochemical analysis of FLT-1 in human Placenta tissue using FLT-1 antibody (Clone: ABM4A56) at 5 µg/ml.