

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-4102: Monoclonal Antibody to RIG-I (Clone: ABM40B5)

Clone Name: Monoclonal
Clone Name: ABM40B5
Application: IHC,FACS,WB

Reactivity: Human
Gene: DDX58
Gene ID: 23586
Uniprot ID: 095786
Format: Purified
Alternative Name: DDX58

**Isotype:** Mouse IgG1 Kappa

Immunogen Information: A partial length recombinant human RIG-I protein (amino acids 1-220) was used as the

immunogen for this antibody.

## **Description**

RIG-I or Probable ATP-dependent RNA helicase DDX58 or simply DDX58 is a RNA helicase enzyme with two isoforms (106 and 101 kDa). RIG-I belongs to the DEAD/H box family and is involved in several cellular responses including RNA binding and alteration of RNA secondary structure. RIG-1 acts as sensor of viral nucleic acids thus activating a cascade of antiviral responses including the induction of type I interferons and pro-inflammatory cytokines. Generally it remains as a monomer but upon activation by viral dsRNA it homomultimerizes and interacts with mitochondria antiviral signaling protein (MAVS/IPS1) which activating a cascade of reactions to produce specific interferon. RIG-1 is mostly expressed in vascular smooth cells.

## **Product Info**

**Amount :**  $25 \mu g / 100 \mu g$ 

**Purification:** Protein G Chromatography

**Content:** 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 2-4  $\mu$ g/ml, Immunohistochemical analysis: 5  $\mu$ g/ml, Flowcytometric analysis: 0.5-1 $\mu$ g/10^6 cells



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-1: Western blot analysis of RIG-I. Anti- RIG-I antibody (Clone: ABM40B5) was used at 2  $\mu$ g/ml on K562 lysates.




Fig-2 : Immunohistochemical analysis of RIG-I in human prostate tissue using RIG-I antibody (Clone: ABM40B5) at 5  $\mu$ g/ml.




Figure-3: Intracellular flow cytometric analysis of RIG-I in K562 Cell line using 0.5  $\mu g/10^6$  cells of Anti-RIGI antibody (ABM40B5). Green represent isotype control and red represent Anti-RIG I antibody (10-4102 Abeomics). Goat anti-mouse PE conjugate was used as secondary.