

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1213: Monoclonal Antibody to Histone H1 (Nuclear Marker)(Clone: HH1/957)

Clonality: Monoclonal HH1/957 Clone Name:

Application: FACS,WB,IF,IHC Reactivity: Human, Mouse, Rat

Gene: H1 3005 Gene ID: **Uniprot ID:** Multiple Format: Purified

Isotype: Mouse IgG2a, kappa

Immunogen Information: Recombinant full-length human Histone H1 protein

Description

Eukaryotic histones are basic and water-soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form the octamer; formed of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. Over 80% of nucleosomes contain the linker Histone H1, derived from an intronless gene that interacts with linker DNA between nucleosomes and mediates compaction into higher order chromatin. Histones are subject to posttranslational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Such modifications include methylation, citrullination, acetylation, phosphorylation, sumoylation, ubiquitination and ADP-ribosylation.

Product Info

Amount: 100 µg

Purification: Affinity Chromatography


100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly Content:

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

Application Note

Flow Cytometry (1-2ug/million cells); Western Blot (1-2ug/ml); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded Rat Pancreas stained with Histone H1 Monoclonal Antibody (HH1/957)




9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982


Email: info@abeomics.com



Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with Histone H1 Monoclonal Antibody (HH1/957)



Formalin-fixed, paraffin-embedded human Tonsil stained with Histone H1 Monoclonal Antibody (HH1/957)



Formalin-fixed, paraffin-embedded human Tonsil stained with Histone H1 Monoclonal Antibody (HH1/957)