

10-4120: Monoclonal Antibody to CLEC4E (Clone: ABM42C6)

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | ABM42C6 |
| Application : | FACS,WB |
| Reactivity : | Human |
| Gene : | CLEC4E |
| Gene ID : | 26253 |
| Uniprot ID : | Q9ULY5 |
| Format : | Purified |
| Alternative Name : | CLEC4E,CLECSF9,MINCLE,UNQ218/PRO244 |
| Isotype : | Mouse IgG1 Kappa |
| Immunogen Information : | A full length human CLEC4E protein was used as the immunogen for this antibody. |

Description

CLEC4E (C-type lectin domain family 4, member E), also called as Mincle, is a type II transmembrane C-type lectin receptor that specifically binds to the cord factor TDM (trehalose-6,6-dimycolate), a unique glycolipid of mycobacterial cell-surface components, and activates immune cells to confer adjuvant activity. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signaling, glycoprotein turnover, and roles in inflammation and immune response. CLEC4E is mainly expressed in myeloid cells, such as macrophages and neutrophils and possesses a single carbohydrate-recognition domain (CRD) in the extracellular region. It is associated with FcRgamma chain, an immunoreceptor tyrosine-based activation motif (ITAM)-containing adaptor, and acts as an activating receptor that senses damaged cells. CLEC4E is also a downstream target of CCAAT/enhancer binding protein (C/EBP), beta (CEBPbeta) and may play a role in inflammation.

Product Info

| | |
|----------------------------|---|
| Amount : | 25 µg / 100 µg |
| Purification : | Protein G Chromatography |
| Content : | 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

Western blot analysis: 2-4 µg/ml, Flowcytometric analysis: 0.5-1 µg/10⁶ cells




Figure-1: Western blot analysis of CLEC4E. Anti-CLEC4E antibody (Clone: ABM42C6) was tested at 2 µg/ml on THP-1 lysate.




Figure-2: cell surface flowcytometric analysis of Anti-CLEC4E in A431 cells using 0.5 µg/10⁶ cells of Anti-CLEC4E (clone: ABM42C6) . Green represent isotype control and red represent Anti-CLEC4E antibody (10-4120 ABEOMICS). Goat anti-mouse PE conjugate was used as the secondary antibody.




Figure-3: cell surface flowcytometric analysis of Anti-CLEC4E in HeLa cells using 0.5 µg/10⁶ cells of Anti-CLEC4E (clone: ABM42C6) . Green represent isotype control and red represent Anti-CLEC4E antibody (10-4120 ABEOMICS). Goat anti-mouse PE conjugate was used as the secondary antibody.